

XV. NOTES ON CRUSTACEA DECAPODA
IN THE INDIAN MUSEUM.

X. HYMENOSOMATIDAE.

By STANLEY KEMP, B.A., *Superintendent, Zoological Survey
of India.*

The small crabs belonging to the family Hymenosomatidae are singularly unobtrusive in habit and unless very abundant are liable to escape notice. A few species are found in salt water of no great depth, and are not infrequently taken on coral reefs or living under stones between tide-marks; but the majority (at any rate on the Indian coast) appear to inhabit estuaries or lagoons where the water is of low or variable salinity. Two species of the family have, indeed, succeeded in establishing themselves in pure fresh water and one has been taken in lakes 3,000 ft. above sea level.¹

Most of the species prefer a bottom composed of mud, which, when matted with the fine hairs on their bodies, doubtless assists them in escaping detection. In many instances the mud forms such a dense coating on the carapace and appendages that it is almost impossible to remove it without injury to the specimen. The legs are very brittle; some species appear to throw them off almost without provocation, and this so constantly occurs with *Elamena* (*Trigonoplax*) *unguiformis* that it is almost impossible to preserve a perfect example.

Among the crabs recently collected on the Indian coasts several species of Hymenosomatidae are represented. Alcock in his memoir on the Indian Catometopes² was able to give an account of five species and two more have since been recorded. Six others, all of which have not hitherto been described, are here added, bringing the total number of known Indian forms up to thirteen.

The new Indian species were all obtained in brackish water. Four were found by myself in Portuguese India, one being a very abundant species which has also been collected by Dr. F. H. Gravely in the Cochin backwaters. The other two were taken by Dr. Annandale and myself in the vicinity of Calcutta. Both these species exhibit very peculiar structure and one of them, obtained on the banks of the River Hughli, cannot be included in any of the genera hitherto described. There can be no doubt that numbers

¹ *Halicarcinus lacustris* (Chilton) [see p. 247, footnote] and *Rhynchoplax introversus*, sp. nov.

² Alcock, *Journ. Asiat. Soc. Bengal*, L.XIX, p. 385 (1900).

of additional species yet remain to be discovered on the Indian coasts.

I have included descriptions of two new forms obtained by Dr. Annandale during his recent tour in the Far East. One of these is from the Tai Hu in the Kiangsu province of China, a lake which is fresh at all times of the year; the other was found in brackish water in the Tale Sap in Lower Siam.

We are at present very far from possessing a clear knowledge of the species referred to this family. The descriptions and figures of many of the older authors are a constant source of difficulty and the identity of numerous species described in the earlier half of the nineteenth century still remains obscure. The confusion is accentuated by differences of opinion regarding the genera. Many authors appear to have distributed their species almost at random and Haswell,¹ who places all the Australian forms in the genus *Hymenosoma*, has expressed the opinion that "the subdivision. . .




FIG. 1.—*Hymenosoma orbiculare*, Desmarest.
Anterior part of carapace, seen from below.

into the genera *Hymenosoma*, *Hymenicus* and *Halicarcinus* appears to be unnecessary and based on extremely slight points of distinction." This view finds no support from subsequent writers, and it is evident that its author was unaware of the characters of the true *Hymenosoma*; nevertheless, as explained below, I believe him to have been right in uniting *Hymenicus* and *Halicarcinus*.

The following notes on the genera are based on the material in the Indian Museum, which contains in addition to twelve Indian species, a number of specimens from China, Australia, New Zealand, S. Africa and the Falkland Is., all the known genera with one exception being represented.

Hymenosoma was described by Desmarest in 1825,² the type species being *H. orbiculare* from the Cape of Good Hope. It is one of the most clearly defined of the genera comprised in the family, differing widely from all others in the complete absence of the epistome (text-fig. 1). The external maxillipedes almost encroach

¹ Haswell, *Cat. Australian Crust.*, p. 114 (1882).

² Desmarest, *Consid. gén. Crust.*, Paris, p. 163 (1825).

on the bases of the antennules and the buccal cavern is not limited anteriorly by a ridge. The ischium of the external maxillipedes is a little longer than the merus; both segments are slender and, when normally folded, gape in the middle line, the underlying appendages being partially visible. In the abdomen of the male the sutures of all the segments are distinct. The regions of the carapace, as in most genera of the family, are defined by fine-cut grooves.

Numerous species have from time to time been placed in *Hymenosoma*, but in the majority of instances the reference is erroneous and it is now practically certain that the genus is monotypic. Stimpson's *H. geometricum*¹ is synonymous with *H. orbiculare* and Guérin Méneville's *H. gaudichaudii*,² though included in the genus by Milne-Edwards,³ is evidently a species of *Halicarcinus*.

Halicarcinus was established by White in 1846,⁴ the type




FIG. 2.—*Halicarcinus planatus* (Fabr.).
Anterior part of carapace, seen from below.

species being Fabricius' *Leucosia planata*⁵ from Tierra del Fuego. In this genus the epistome is a conspicuous plate, and the buccal cavern is bounded anteriorly by a transverse ridge (text-fig. 2). The ischium and merus of the external maxillipedes are of similar size and are broad segments, completely or almost completely closing the buccal cavern. As in *Hymenosoma* the grooves on the upper surface of the carapace are clean cut and, in the abdomen

¹ Stimpson, *Proc. Acad. Sci. Philadelphia*, X, p. 108 [54] (1858) and *Smiths. Misc. Coll.*, XLIX, p. 144 (1907). Stebbing, in *Marine Invest. S. Africa*, IV, p. 50 (1905) and *Ann. S. African Mus.*, VI, p. 332 (1910), retains *H. geometricum* as a distinct species, but has since agreed that it is synonymous with *H. orbiculare* [see *Trans. Roy. Soc. Edinburgh*, L, ii, p. 270 (1914)].

² Guérin Méneville, *Voy. de la 'Coquille'*, II, ii, 1^{re} div., p. 21 and *Atlas, Crust.*, pl. ii, figs. 12-18.

³ Milne-Edwards, *Ann. Sci. nat., Zool., Paris* (3), XX, p. 222 (1853).

⁴ White, *Ann. Mag. Nat. Hist.* (1), XVIII, p. 178 (1846).

⁵ For references see Stebbing, *Proc. Zool. Soc. London*, 1900, p. 524; Doflein and Balss, *Mitth. naturhist. Mus. Hamburg*, XXIX, p. 35 (1912) and Chilton, *Subantarctic Is. of New Zealand*, p. 609 (1910).

of the male, the sutures of all the segments are distinct. In my interpretation of its limits, *Halicarcinus* comprises species with simple rostra as well as those in which it is trilobate or tridentate.

Lucas's *Hombronia*,¹ suggested as a generic name for Jacquinot's *Hymenosoma depressa*² from the Auckland Is. and Nicolet's *Liriopea*,³ based on two species from Chili, are generally regarded as synonyms of *Halicarcinus*, and Dana's *Hymenicus*⁴ is separated by such slight distinctions that it cannot in my opinion be retained as a separate genus. In describing *Hymenicus* Dana says: "In this genus the front has not the three teeth of *Halicarcinus* (between which the flexed first antennae are seen), but a simple rounded or trilobate prominence forms the front, and the first antennae are covered. The feet are much longer and more slender than in any of the species of *Halicarcinus*, seen by the author." On comparing *H. varius*, the type species of *Hymenicus*, with *Halicarcinus planatus*, the points to which Dana has drawn attention are readily appreciated. The difference, however, is in reality of very slight morphological importance and is entirely due to the greater development of the front in *H. varius*, the disposition and structure of the related parts being as nearly as possible identical. Examination of allied forms shows that a wide variation exists in the form of the front and affords conclusive evidence that the character is of specific rather than generic value. The comparatively great length of the legs in *H. varius*—the only other point mentioned by Dana—is clearly insufficient as a generic criterion; the external maxillipedes are almost identical in structure with those of *H. planatus* and, as in that species, the sutures of all the segments of the male abdomen are distinct.

But though Dana's *Hymenicus* must, through the characters of its type species, be placed in the synonymy of *Halicarcinus*, it does not follow that all the species hitherto referred to *Hymenicus* must be transferred to White's genus. The two Indian species described by Alcock,⁵ together with four others dealt with below, appear to offer distinctive characters. In most particulars they agree with *Halicarcinus*, but the external maxillipedes are much more slender, with the ischium conspicuously smaller than the merus; when normally folded they gape widely in the middle line, leaving parts of the underlying appendages exposed (see text-fig. 7, p. 259). In the abdomen of the male, moreover, the 3rd, 4th and 5th segments are fused, with complete obliteration of the sutures (see text-fig. 9, p. 259). The rostrum is variable in form, but is normally tridentate or trilobate.

¹ Lucas, in Hombron and Jacquinot's *Voy. au Pôle Sud, Zool.*, III, Crust. p. 62 (1853).

² Jacquinot, Atlas to above, *Crust.*, pl. v, figs. 34-39 (1842-53); Chilton, *Ann. Mag. Nat. Hist.* (7), XIX, p. 146, pl. v (1907). It is perhaps doubtful whether this species really belongs to *Halicarcinus* as here defined, for the grooves on the upper surface of the carapace are not shown in either of the figures.

³ Nicolet, in Gay's *Hist. física y política de Chile, Zool.*, III, p. 158 (1849).

⁴ Dana, *U. S. Explor. Exped., Crust.*, I, p. 387 (1852).

⁵ Alcock, *Journ. Asiat. Soc. Bengal*, I, XIX, p. 388 (1900).

As a generic name for this group of species I have employed Stimpson's *Rhynchoflax*,¹ though unfortunately I cannot be altogether certain that its application is correct. Stimpson does not state that any parts of the male abdomen are fused and his reference to the external maxillipedes is decidedly confusing, his only remark being "ischium-joint. . . scarcely longer than meros," a description that applies if anything better to *Haliscarcinus* than to the group of Indian species. On the other hand *Rhynchoflax messor*, the type species of the genus from Simoda in Japan, appears specifically to be an exceedingly close relative of Alcock's "*Hymenicus*" *wood-masoni*, both species, apart from other resemblances, possessing a series of teeth on the upper border of the merus of the chelipede. The question cannot finally be settled until further specimens of *Rhynchoflax messor* are examined. The types were, I understand, destroyed by fire in 1871 and the species has not been recorded since Stimpson's time.

To distribute the numerous described species correctly between the genera *Haliscarcinus* and *Rhynchoflax*, as here defined, is a matter of very great difficulty, but from the figures and descriptions which have been published I conclude that the following species may safely be referred to the genus *Haliscarcinus*,²—*Hymenosoma gaudichaudii*, Guérin Méneville,³ *Haliscarcinus pubescens*, Dana,⁴ *Hymenicus pubescens*, Dana,⁵ *Hymenicus varius*, Dana,⁶ *Haliscarcinus ovatus*, Stimpson,⁷ *Hymenosoma tridentata*, Jacquinot,⁸ *Hymenosoma rostratum*, Haswell,⁹ *Elamene pilosa*, A. Milne-Edwards,¹⁰ *Hymenosoma laeue*, Targioni-Tozzetti,¹¹ *Hymenicus marmoratus*, Chilton,¹² and *Hymenosoma lacustris*, Chilton.¹³

¹ Stimpson, *Proc. Acad. Sci. Philadelphia*, X, p. 109 [55] (1858) and *Smiths. Misc. Coll.*, XLIX, p. 147 (1907).

² In addition to the type species of the genus I have seen specimens of *H. ovatus*, *H. varius*, *H. rostratus* and a species from the Australian coast which is perhaps undescribed.

³ Guérin Méneville, *loc. cit. supra* p. 245.

⁴ Dana, *U. S. Explor. Exped., Crust.*, I, p. 386, pl. xxiv, fig. 8.

⁵ Dana, *ibid.*, p. 388, pl. xxiv, figs. 11 a-c.

⁶ Dana, *ibid.*, p. 387, pl. xxiv, fig. 9.

⁷ Stimpson, *Proc. Acad. Nat. Sci. Philadelphia*, X, p. 109 [55] (1858) and *Smiths. Misc. Coll.*, XLIX, p. 146 (1907); Stebbing, *Proc. Zool. Soc. London*, 1900, p. 525, pl. xxxvi. Chilton, in *Subantarctic Is. New Zealand*, p. 609 (1910) suggests that *H. ovatus* is synonymous with Jacquinot's *H. tridentata*.

⁸ Jacquinot, in Hombron and Jacquinot's *Voy. au Pôle Sud, Zool., Atlas. Crust.*, pl. v, figs. 27-33. Usually regarded as a synonym of *H. planatus*. Chilton, *loc. cit.*, 1910, p. 609, suggests its retention at least in a subspecific significance.

⁹ Haswell, *Proc. Linn. Soc. N. S. Wales*, VI, p. 550 (1882) and *Cat. Australian Crust.*, p. 116 (1882); Baker, *Trans. Roy. Soc. S. Australia*, XXX, p. 114, pl. iii, figs. 2, 2a,b (1906).

¹⁰ A. Milne-Edwards, *Nouv. Arch. Mus. Paris*, IX, p. 322, pl. xviii, figs. 6, 6a-e (1873).

¹¹ Targioni Tozzetti, *Crost. Viaggio 'Magenta'*, p. 179, pl. xi, figs. 3a-e (1877).

¹² Chilton, *Trans. N. Zealand Inst.*, XIV, p. 172, pl. viii, figs. 1a-c (1881).

¹³ Chilton, *Trans. N. Zealand Inst.*, XIV, p. 172 (1881) [as *Elamene? lacustris*]; *ibid.*, XLIV, p. 128 (1912); *ibid.*, XLVII, p. 316, fig. 1 (1915); Fulton and Grant, *Proc. Roy. Soc. Victoria*, XV, p. 59, pl. viii (1902); Grant and McCulloch, *Proc. Linn. Soc. N. S. Wales*, XXXII, p. 153 (1907).

The generic position of a number of other species is doubtful, but I think it will eventually be found that all those from southern latitudes hitherto referred to the genera *Hymenicus* and *Hymenosoma*¹ belong in reality to *Halicarcinus*. *H. planatus*, if the records are to be trusted, is circumpolar in distribution and the species listed above are without exception from southern latitudes. The forms that can be referred to *Rhynchoplax* are, on the other hand, all found on the Asiatic coasts, from which no representative of *Halicarcinus* has yet been obtained.

The genus *Rhynchoplax*, in my estimation, comprises Stimpson's two species, *R. messor* from Japan and *R. setirostris* from Hong Kong, de Man's *Elamene filholi*² from near Batavia, Alcock's *Hymenicus wood-masoni* and *H. inachoides* from India and six other species described below. It probably includes also Miss Rathbun's *R. coralicola*³ from Singapore.

A species of Hymenosomatidae found on the banks of the R. Hughli, near Calcutta, does not appear to be admissible into any of the genera hitherto recognized; it is described below under the name *Hymenicoides carteri*. In its structure this species shows a high degree of specialization and generically is related to *Halicarcinus* and *Rhynchoplax*. It agrees with the former of these genera in having the sutures of all the segments of the male abdomen distinct and with the latter in the slender form of the basal segments of the third maxillipedes: it differs from both in the remarkably elongate dactylus of the latter appendages and in the entire absence of a rostrum (see text-fig. 16, p. 267).

In *Hymenicoides* the antennule is completely exposed in dorsal view. This character has frequently been used as a generic criterion, but in my opinion is of specific importance only, being due almost entirely to the extent to which the rostrum is reduced. In *Rhynchoplax* the rostrum is normally trilobate and well developed, but in *R. nasalis*, sp. nov., the lateral portions are suppressed, with the result that the antennules, just as in *Hymenicoides carteri*, are visible from above.

The genus *Elamena* was established by Milne-Edwards in 1837,⁴ the type species being Desmarest's *Hymenosoma mathaei*⁵ from the Ile de France. Haswell's suggestion that this species is merely the young of *Halicarcinus planatus*⁶ has been contested by Stebbing and is certainly incorrect. That Rüppell's identification⁷ of Desmarest's species is correct may be assumed from the

¹ Except, of course, *Hymenosoma orbiculare* and the synonymous *H. geometricum*.

² De Man, *Arch. f. Naturgesch.*, LIII, i, p. 386, pl. xvii, fig. 3 (1887).

³ Rathbun, *K. Danske Vid. Selsk. Skrift.* (7), *naturvid. og math.*, V, p. 316, text-fig. 5 (1910).

⁴ Milne-Edwards, *Hist. nat. Crust.*, II, p. 33 (1837).

⁵ Desmarest, *Consid. gén. Crust.*, Paris, p. 163 (1825). I have not seen this species.

⁶ Haswell, *Cat. Australian Crust.*, p. 114 (1882).

⁷ Rüppell, *Beschreib. Abbild. 24 Arten Krabben*, Frankfurt, p. 21, pl. v, fig. 1 (1830).

fact that his description and figure is quoted by Milne-Edwards, who, as Stebbing has remarked, probably had Desmarest's specimen before him when he wrote. Paulson's figure¹ differs considerably from that given by Rüppell; the carapace is of much greater proportionate length and bears grooves on its upper surface much as in *Halicarcinus* and *Rhynchoplax*. Stebbing has pointed out (*loc. cit.*) that Milne-Edwards' subsequent reference to the species in 1853² is almost certainly erroneous; the genus is here credited with a tridentate rostrum, a character not found in Desmarest's species.

In *Elamena*, as represented by the species in the Indian Museum, the carapace is very greatly depressed, sometimes of wafer-like thinness, and the regions of its upper surface are not defined by the fine-cut grooves found in the other genera. The epistome is conspicuous and sometimes of great length. The external maxillipedes completely close the buccal cavern and the ischium, though somewhat variable in size, is always longer than the merus (see text-fig. 25, p. 276). As in *Rhynchoplax* the 3rd, 4th and 5th segments of the male abdomen are fused, and the sutures between them obliterated. The front, or rostrum, is simple, never trilobate.

I agree with Alcock³ that *Trigonoplax* is, at most, only a subgenus of *Elamena*. It was described by Milne-Edwards in 1853,⁴ the type species being de Haan's *E. unguiformis*.⁵ As has been pointed out above, Milne-Edwards when writing in 1853 appears to have misunderstood the characters of his own genus *Elamena*, and the foundation of *Trigonoplax* seems to have been a direct result of this mistake. The only constant differences that I am able to find between *Elamena* and *Trigonoplax* do not appear to be important and it is probable that when the characters of the species are better understood, the latter will come to be regarded as a synonym of the former.

Six species of *Elamena* have been found on the Indian coasts and are referred to below; of these three (perhaps four) belong to the subgenus *Trigonoplax*. Other representatives of the genus are *E. mathaei* (Desmarest),⁶ the type species, found at Réunion and in the Red Sea, *E. producta*, Kirk⁷ (with which *E. kirki*, Filhol,³ is

¹ Paulson, *Crust. Red Sea, Kiew*, p. 71, pl. ix, figs. 3, 3a,b (1871).

² Milne-Edwards, *Ann. Sci. nat., Zool., Paris* (3), XX, p. 223, pl. xi, figs. 4, 4a (1853).

³ Alcock, *Journ. Asiat. Soc. Bengal*, l. XIX, p. 386 (1900).

⁴ Milne-Edwards, *Ann. Sci. nat., Zool., Paris* (3), XX, p. 224 (1853).

⁵ De Haan, in Siebold's *Fauna Japonica. Crust.*, p. 75, pl. xxix, fig. 1, pl. H (1839).

⁶ For references see p. 248, footnotes 4, 5. *Elamene truncata*, Lenz (not A. M.-Edw.), *Abhandl. Senckenberg. Naturforsch. Ges. Frankfurt*, XXVII, i, p. 367, pl. xlviii, figs. 15a,b (1902) is apparently synonymous.

⁷ Kirk, *Trans. N. Zealand Inst.*, XI, p. 395 (1878); Filhol, *Recueil de Mém. Inst. France, Miss. à l'île Campbell, Zool.*, p. 404, pl. 1, figs. 1, 2 (1885); Chilton, *Rec. Canterbury Mus.*, I, p. 294 (1911).

⁸ Filhol, *loc. cit. supra*, p. 405, pl. xlvii, figs. 6, 8 (1885).

apparently synonymous) and *E. longirostris*, Filhol,¹ both from New Zealand. The position of *E. quoyi*, Milne-Edwards,² *E. mexicana*, Milne-Edwards³ and *E. whitei*, Miers,⁴ is doubtful. *E. pilosa*, A. Milne-Edwards, as already pointed out, is probably a species of *Halicarcinus*, while *E. filholi*, de Man, appears to belong to *Rhynchoplax*. *E. minuta*, A. Milne-Edwards,⁵ whatever it may be, is certainly not an *Elamena*.

Elamenopsis was established by A. Milne-Edwards in 1873 for *E. lineatus*,⁶ a species found in New Caledonia. I have seen no specimens of the genus and have not been able to satisfy myself regarding its position in the family. It is said to form a link between the Hymenosomatidae and Pinnotheridae. From the description it appears to be related to *Rhynchoplax*, but the walking legs are much shorter and stouter than in any species of that genus that I have seen.

The principal characters of the other five genera may be summarised in the following way:—

- I. There is no epistome. [The external maxillipedes are slender and do not nearly close the buccal cavern. In the abdomen of the male the sutures of all the segments are distinct] *Hymenosoma*,
Desmarest.
- II. The epistome is well defined and frequently very long.
 - A. The regions of the carapace are defined by sharp-cut grooves. The ischium of the external maxillipedes is not longer, frequently much shorter than the merus.
 1. A rostrum is present and is frequently trilobate or tridentate. The dactylus of the external maxillipedes is short (normal).
 - a. The external maxillipedes are broad and completely, or almost completely, close the buccal cavern. In the abdomen of the male the sutures of all the segments are distinct *Halicarcinus*,
White (= *Hymenicus*, Dana).
 - b. The external maxillipedes are slender and do not nearly close the buccal cavern. The 3rd, 4th and 5th segments of the male abdomen are fused and the sutures obliterated *Rhynchoplax*,
Stimpson.
 2. The rostrum is altogether absent. The dactylus of the external maxillipedes is abnormally long, reaching the hinder limit of the buccal cavern. [The external maxillipedes are very slender and do not nearly close the buccal cavern. In the abdomen of the male the sutures of all the segments are distinct]... *Hymenicoides*,
gen. nov.

¹ Filhol, *loc. cit. supra*, p. 403, pl. xlvi, fig. 7 (1885).

² Milne-Edwards, *Ann. Sci. nat., Zool., Paris* (3), XX, p. 223, pl. xi, fig. 3 (1853).

³ Milne-Edwards, *ibid.*, p. 224.

⁴ Miers, *Cat. Crust. N. Zealand*, p. 52, pl. i, fig. 4 (1876).

⁵ A. Milne-Edwards, *Nouv. Arch. Mus. Paris*, IX, p. 324, pl. xviii, fig. 5 (1873).

⁶ A. Milne-Edwards, *ibid.*, p. 324, pl. xviii, fig. 4.

B. The surface of the carapace is smooth, rarely uneven; its regions are never delimited by sharp-cut grooves. The ischium of the external maxillipedes is longer than the merus. [The rostrum, when present, is simple. The external maxillipedes are broad and completely close the buccal cavern. The 3rd, 4th and 5th segments of the abdomen of the male are fused and the sutures obliterated]

Elamena, Milne-Edwards.

Of these genera only the last three are found on the Indian coasts.

Genus *Rhynchoplax*, Stimpson.

1858. *Rhynchoplax*, Stimpson, *Proc. Acad. Sci. Philadelphia*, X, p. 100 [55].

1900. *Hymenicus*, Alcock (not of Dana), *Journ. Asiat. Soc. Bengal*, LXIX, p. 387.

1907. *Rhynchoplax*, Stimpson, *Smiths. Misc. Coll.*, XLIX, p. 147.

The carapace is circular, ovate or polygonal in outline and is depressed; the upper surface is sunken with the usual grooves sharply defined and the margin upturned. The rostrum is tridentate or trilobate, the lateral processes very rarely absent. The epistome is of good length and the buccal cavern is bounded anteriorly by a sharp ridge. The external maxillipedes are comparatively slender and, when normally folded, gape widely in the middle line; the merus is longer than the ischium and the dactylus is, as usual, short. The chelipedes in both sexes are stouter than the walking legs. In the abdomen of the male the 3rd, 4th and 5th segments are fused and the sutures between them completely obliterated.

This genus is very closely related to *Halicarcinus*, but is distinguished by the more slender merus and ischium of the external maxillipedes and by the fact that certain segments of the male abdomen are fused.

The six Indian species of *Rhynchoplax* together with the two obtained by Dr. Annandale in Siam and China may be distinguished in the following manner:—

1. A large forwardly directed tooth or process on either side of carapace above base of 1st walking legs [rostrum tridentate; a sharp post-ocular tooth visible in dorsal view].

A. Carapace subcircular, its antero-lateral border armed with one or two blunt teeth; merus of chelipede armed with several strong teeth on its upper border; dactyli of last three legs armed with a series of small teeth.

1. Two teeth on antero-lateral border of carapace; chela of adult male more than twice as long as high, palm rounded below, fingers not gaping and armed with regular teeth

R. wood-masoni
(Alcock).

2. Only one tooth on antero-lateral border of carapace; chela of adult male much less than twice as long as high, palm keeled below, fingers widely gaping and with irregularly disposed teeth

R. alcocki, sp.
nov.

- B.* Carapace octagonal; its antero-lateral border without teeth; merus of chelipede without teeth on its upper border; dactyli of last three legs with a single large tooth near apex *R. octagonalis*,
sp. nov.
- II. No tooth on side of carapace above base of 1st walking legs [no teeth on antero-lateral border of carapace].
- A.* Rostrum trilobate or tridentate; basal segment of antennular peduncle not visible in dorsal view; penultimate piece of abdomen of male longer than broad, without tubercle.
1. Rostrum composed of three very broad lobes; post-ocular tooth not visible from above; legs stout, dactyli without teeth *R. demeloi*,
sp. nov.
2. Rostrum composed of three narrow lobes or teeth; post-ocular tooth visible from above; legs slender, dactyli with teeth.
- a.* Carapace not longer than broad; rostrum composed of three lobes; 2nd walking legs not more than $2\frac{1}{2}$ times length of carapace.
- i. Postero-lateral border of carapace normal; dactyli of walking legs very strongly curved and with very large teeth; terminal segment of male abdomen broader than long *R. exiguus*,
sp. nov.
- ii. Side-walls of branchial region of carapace reflected upwards, forming a crest outside the true postero-lateral border; dactyli of walking legs moderately curved, with small teeth; terminal segment of male abdomen much longer than broad *R. introversus*,
sp. nov.
- b.* Carapace much longer than broad, rostrum composed of three long teeth; 2nd walking legs more than 3 times length of carapace [dactyli of walking legs moderately curved, with small teeth; terminal segment of male abdomen as long as broad] *R. inachoides*,
Alcock.
- B.* Rostrum composed of a single tooth-like process; basal segment of antennular peduncle completely visible in dorsal view; penultimate piece of abdomen of male broader than long, with a large tubercle at distal end. [no post-ocular tooth; dactyli of last three legs with a single tooth] *R. nasalis*,
sp. nov.

Rhynchoplax wood-masoni (Alcock).

1900. *Hymenicus wood-masoni*, Alcock, *Journ. Asiat. Soc. Bengal*, LXIX, p. 388, and (1902) *Illust. Zool. 'Investigator'*, pl. lxiv, fig. 4.

A few particulars regarding the structure of this species, some of which are additions to Alcock's description, are given below in the course of a comparison with the closely allied *R. alcocki*.

Stimpson's *Rhynchoplax messor* from Simoda appears also to be a related form, agreeing in the presence of a series of teeth on the upper aspect of the carpus of the chelipede. In the Japanese

species, however, the carapace is stated to be triangular, with only two teeth on the lateral border, and the median tooth of the rostrum points obliquely upwards, instead of being depressed as in *R. wood-masoni*. The carpus of the chelipedes bears on its upper surface three or four small teeth; in adult males of *R. wood-masoni* one such tooth is sometimes found near the meral articulation, but it is frequently absent. Stimpson states that each joint of the ambulatory feet, except the dactyli, is "dentigerous in the middle," a character not found in *R. wood-masoni* or in any other species of the genus that I have seen.

The only specimens of this species in the Indian Museum are those described by Alcock from the Andamans and from Port Canning near Calcutta.




FIG. 3.—*Rhynchoplax alcocki*, sp. nov.

***Rhynchoplax alcocki*, sp. nov.**

The carapace is subcircular, a little produced anteriorly and with its sides slightly flattened and nearly parallel. Its breadth is almost equal to its length, excluding the rostrum. The surface is hairy and sunken and the usual grooves are well defined. The entire margin is upturned and is continuous from side to side across the base of the rostrum. A sharp post-ocular tooth is visible in dorsal view and behind it, on the margin itself, there is a blunt tooth corresponding to the foremost of those found in *R. wood-masoni* (text-fig. 4*d*). Below the margin near the base of the first pair of walking legs there is a huge tooth-like process directed forwards, upwards and outwards.

The rostrum is composed of three narrow lobes with rounded extremities; the median lobe is longer than the two others, and its apex is situated on a lower level.

The antennules, when folded, are concealed beneath the rostrum; at their base they are separated by a prominent septum. The epistome is of moderate length. As in *R. wood-masoni* the external maxillipedes are slender and do not nearly close the




FIG. 4.—*a-c*, *Rhynchoplax wood-masoni* (Alcock).
d-g, *Rhynchoplax alcocki*, sp. nov.

a, d.—Rostrum, eye and antero-lateral margin of carapace.

b, e.—Chela of male (denuded).

c, f.—Abdomen of male.

g.—Terminal part of dactylus of penultimate leg.

buccal cavern. The ischium is produced at its inner distal angle and the merus is expanded antero-externally, partially concealing the exognath.

The chelipedes in both sexes are stouter than the walking legs, the chelae of the adult male being particularly large. In the male the merus bears a conical tooth near the end of its lower margin and a series of some five large blunt teeth superiorly.

The carpus is smooth. The chela is very greatly compressed and in adult males is little more than one and a half times as long as high. The palm in lateral view is nearly circular in outline (text-fig. 4e) and is slightly hollowed both internally and externally near the strongly compressed upper border. The lower border is convex and is keeled. The fingers gape very widely and meet only at their tips. The dactylus is almost twice the length of the upper border of the palm and bears in its basal third a single large tooth, in front of which a second smaller tooth is occasionally found. The fixed finger bears two large teeth in advance of those on the dactylus and one or two others, which are smaller, near the base. Near the apex, where they meet, the fingers are provided with four or five small interlocking teeth. In the female the teeth on the merus are obscure or altogether wanting and the chela is much narrower, fully twice as long as high; the fingers meet throughout their length and are armed with regularly spaced teeth.

The walking legs are very slender; those of the second pair are slightly the longest and are about two and a quarter times the length of the carapace and rostrum. The anterior border of the merus ends in a very obscure tooth. The dactyli are very slender and are curved; close to the apex each is armed with a large recurved tooth (text-fig. 4g) and in front of this, in the last three pairs, there is a series of 8 to 11 smaller teeth, also recurved and extending over practically the whole length of the posterior margin. The chelipedes and legs are clothed with hair, which is particularly long and thick on the chela of the male.

The sternum and abdomen are densely clothed with hair. The abdomen of the male resembles that of *R. wood-masoni*, but is slightly narrower. The terminal segment is scarcely longer than broad and is rounded at the apex (text-fig. 4f); the preceding portion is longer than broad, parallel-sided at the base and from the middle point onwards strongly narrowed.

A large male is only 4.8 mm. in length from the tip of the rostrum to the posterior margin of the carapace. Ovigerous females are smaller, sometimes not more than 4 mm. long. The carapace of living specimens, when brushed clean, was of a dull purplish brown colour with groups of small whitish spots.

R. alcocki is very closely allied to *R. wood-masoni*, Alcock, there being an almost exact resemblance between the two in the teeth on the merus of the male chelipede. Apart from size, *R. wood-masoni* being much the larger form, the species may be distinguished by the following characters (cf. text-figs. 4a-c and 4d-f):—

R. wood-masoni, Alcock.

Carapace longer, its length excluding rostrum about one-tenth greater than its breadth.

Two teeth on antero-lateral margin of carapace.

R. alcocki, sp. nov.

Carapace shorter, its length excluding rostrum scarcely greater than its breadth.

Only one tooth on antero-lateral margin of carapace.

Rostral teeth slender.
Cornea of eye proportionately smaller.

Chela of adult male not greatly compressed, more than twice as long as high, lower edge of palm rounded.

Fingers of chela of adult male not gaping at base, armed with a regular series of teeth.

Terminal segment of abdomen of male apically pointed.

Rostral teeth less slender.
Cornea of eye proportionately larger.

Chela of adult male very greatly compressed, much less than twice as long as high, lower edge of palm keeled.

Fingers of chela of adult male widely gaping at base, armed with a very irregular series of teeth.

Terminal segment of abdomen of male apically rounded.

There are altogether about 100 specimens of this species in the Indian Museum. The greater number were found in Portuguese India in September 1916 and were obtained in the Rachol river at the head of Mormugao Bay above Cortalim Point and in the Mandavi river at Nova Goa. Some of the specimens were dredged on a muddy bottom in water from $1\frac{1}{2}$ to $4\frac{1}{2}$ fathoms in depth, while others were found at Betim Point opposite Nova Goa, living on the posts of a jetty densely covered with Hydroid. All the specimens were found in brackish water, the specific gravity (corrected) varying from about 1.0010 to 1.0060. There are also in the Museum a few specimens found by Dr. F. H. Gravely in September 1914 in the Cochin backwaters near Ernakulam.

The types are from Portuguese India and bear the number 9735/10 *Zool. Surv. Ind.*

Rhynchoplax octagonalis, sp. nov.

The carapace, rostrum excluded, is a trifle broader than long and is distinctly octagonal in outline. The surface, in an ovigerous female, is very little sunken; it is rather closely covered with short hairs and the usual grooves are well defined. The margin is continuous from side to side across the base of the rostrum and is entire, the blunt teeth found on the antero-lateral borders in the two preceding species being absent. On the side wall above the base of the first pair of walking legs there is a large and sharp procurved tooth (text-fig. 5).

The rostrum in dorsal view is seen to consist of three sharp isolated spines, the lateral ones a little shorter than the median and directed obliquely outwards and upwards. The greater part of the eye can be seen from above, together with a small but sharp post-ocular tooth.

The antennules when folded are not visible in dorsal view; they are separated by a well-marked septum. The external maxillipedes resemble those of the preceding species.

The chelipedes of the female are stouter than the legs. The merus does not bear any distinct teeth. The chela is not com-

pressed and the fingers, which are longer than the palm, meet throughout their length when the claw is closed and bear a regular series of 5 or 6 teeth on their inner margins.

The walking legs are not very slender; those of the second pair are about twice the length of the carapace and rostrum. The anterior border of the merus in each pair ends in a prominent tooth. The dactylus of the first walking legs is unarmed; that of the three following pairs is provided with a stout recurved tooth close to the apex. The chelipedes bear scattered hairs; these also occur on the walking legs, which are, moreover, densely fringed on their posterior margins.




FIG. 5.—*Rhynchoplax octagonalis*, sp. nov.

The species is described from a single ovigerous female, with carapace about 3.9 mm. in length.

De Man's *Elamene filholi*¹, from Noordwachter I. near Batavia, is without doubt a species of *Rhynchoplax* and resembles *R. octagonalis* in the structure of the dactyli of the walking legs and in the position of the single tooth found on the lateral margin of the carapace. In the Javanese species, however, the eye is altogether concealed from above, the carapace and rostrum are quite different in form and the legs are much more slender. Miss Rathbun's *R. coralicola*² from Singapore also possesses a single tooth at the

¹ De Man, *Archiv. f. Naturgesch.*, LIII, i, p. 386, pl. xvii, fig. 3 (1887).

² Rathbun, *K. Danske Vid. Selsk. Skrift.* (7), *naturvid. og math.*, V, p. 310, text-fig. 5 (1910).

side of the carapace, but it is said to be antero-lateral in position. In this species the dactyli are spinulose, thus differing conspicuously from those of *R. filholi* and *R. octagonalis*.

The specimen was obtained at low water under stones among mangroves on Vareeg Islet in Mormugao Bay, Portuguese India. It bears the number 9740/10 *Zool. Surv. Ind.*

***Rhynchoplax demeloi*, sp. nov.**

The carapace is nearly circular; the breadth of its upper surface is about equal to its length, including the median rostral lobe.



FIG. 6.—*Rhynchoplax demeloi*, sp. nov.

The surface is greatly sunken and is covered with fine hairs that retain a quantity of mud; the usual grooves are deeply cut. The lateral border is entire, upturned, and continuous anteriorly across the base of the three rostral prominences; it is obscurely angulate a short distance behind the eye. The tooth found in the three preceding species on the side wall of the carapace is absent.

The three rostral prominences are exceedingly short. The median one is almost square, a little longer than broad, and is abruptly deflexed; the other two are rounded, very much broader than long, and project straight forwards. The greater part of the cornea of the eye is visible in dorsal view.

When viewed from below the median rostral lobe is seen to be longitudinally carinate and behind the eye there is a small post-ocular tooth which is altogether invisible from above (text-fig. 7). The antennules when folded are completely concealed beneath the front; they are separated at their base by a strong septum. The epistome is rather short. The external maxillipedes are similar to those of *R. alcocki*.




FIG. 7.—*Rhynchoplax demeloi*, sp. nov.
Anterior part of carapace, seen from below.

The chelipedes in both sexes are stouter than the walking legs and the chelae are much larger in the male than in the female. The merus and carpus are without teeth. The chela of the adult male (text-fig. 8) is about twice as long as high and is not carinate on its upper or lower margins. Except for a gap close to the base the fingers meet throughout their length; they are armed on their inner margins with 5 or 6 broad interlocking teeth that diminish in size from behind forwards. The dactylus is nearly twice the




FIG. 8.—*Rhynchoplax demeloi*, sp. nov.
Chela of male (denuded).

length of the upper border of the palm. The chela of the female is similar, but more slender. In both sexes the chelipedes are covered with fine hairs; on the outer surface of the palm of the male they are very long and dense, each retaining a quantity of mud.

The second walking legs are slightly the longest and are a little more than two and a half times the length of the carapace. All the segments are exceptionally broad and the anterior border of the merus in each pair ends in a blunt tooth. The dactyli are quite flat, very broad, and only slightly curved; that of the last pair is only about four times as long as wide. The posterior margin is without any of the usual recurved teeth, in this respect differing from all other Indian species of the genus. The walking legs like all other parts of the body are covered with fine hairs which form a short but dense fringe on the posterior borders of the last four segments.



FIG. 9.—*Rhynchoplax demeloi*, sp. nov.
Abdomen of male.

The abdomen of the male is similar to that of *R. alcocki* and *R. wood-masoni*, but is narrower. The ultimate segment is bluntly

pointed and is much longer than broad; the preceding portion, which, as in the other species, appears to comprise three fused segments, is obscurely grooved in the middle line; its lateral margin is angulate near the middle and in front of this point is distinctly concave.

The carapace of the largest specimen, an adult male, is about 4.4 mm. in length.

The species is described from fourteen specimens, including a number of ovigerous females, obtained on the shores of the Mandavi river at Nova Goa in Portuguese India. They were found at low water under stones on a muddy bank. At the time they were taken the water in the river was brackish, the specific gravity being about 1.0010.

With this species I have associated the name of Capt. Froilano de Melo, Director of the Bacteriological Laboratory of the Instituto de Análises e Vacina at Nova Goa. I am greatly indebted to Capt. de Melo for the assistance he gave me during my visit to Portuguese India, especially for facilities for the investigation of the very interesting fauna of the Mandavi river.

The types bear the number 9741/10 *Zool. Surv. Ind.*

Rhynchoplax exiguus, sp. nov.

The carapace is ovate and is widest a little behind the middle point; its upper surface, rostrum included, is a little longer than broad. In an adult female (text-fig. 10) the portions of the carapace above the bases of the first two pairs of walking legs are swollen, covered with stiff hairs and project beyond the upturned lateral margin of the carapace; in males these parts are not visible in dorsal view. There is no tooth or process above the base of the first walking legs or on the antero-lateral margin. The upper surface is a little sunken, covered with fine hairs, and with the usual grooves sharply defined. The rostrum is composed of three lobes set with stiff setae. The median lobe is depressed and longer than the other two; in the adult female it is narrow and parallel-sided, in males broader at the base and triangular.

The eye is unusually large; the entire cornea and a portion of the stalk is visible from above, together with a large and very conspicuous post-ocular tooth. The antennules when folded are completely concealed beneath the front; at their base they are separated by a well-marked septum.

The epistome is comparatively long. The buccal cavern is of the usual form and is not nearly closed by the external maxillipedes. The merus in the latter appendage is a little longer than the ischium and expanded antero-externally, partially concealing the exopod. The stalk of the exopod, as in *R. naso*, is long and projects a little beyond the endopod when the segments are normally flexed.

The chelipedes of male specimens (which are perhaps not full grown) resemble those of the female, the chela being only a little

stouter than the walking legs. The merus is without teeth and the chela, though the palm is somewhat swollen, is comparatively long and slender. The fingers when closed meet throughout their length and are armed from base to apex with a regular series of 5 or 6 teeth.

The walking legs are slender; those of the second pair are about twice the length of the carapace and rostrum. The merus in each pair ends bluntly. The dactylus is long, slender and very strongly curved; the apex is finely pointed and on the posterior margin there are a number of exceptionally large recurved teeth. In the adult female there are 8 or 9 such teeth, distributed




FIG. 10.—*Rhynchoplax exiguus*, sp. nov.

along the whole length of the dactylus; in smaller specimens they are less numerous—sometimes only 3—and occur only in the distal half.

The abdomen of the male is similar to that of *R. demeloi*. The 3rd, 4th and 5th segments form a single piece; the ultimate segment is triangular and a little broader than long.

The species is described from ten specimens, most of which are exceedingly small. The adult female, which appears to have been captured soon after the eggs were discharged, is only 3.4 mm. in length from the tip of the rostrum to the hinder part of the carapace. The largest male, similarly measured, is only 2.8 mm. in length.

Rhynchoplax exiguus appears to be related to *R. inachoides* (Alcock), but is distinguished by the broader carapace, shorter rostral lobes, much shorter walking legs and larger dactylar teeth.

The specimens were obtained by Dr. Annandale in the Tale Sap in Peninsular Siam. They were found on the mainland opposite the western end of Koh Yaw, living in lumps of turf that had fallen into the lake owing to the undermining of the bank. The water in the vicinity was brackish, the specific gravity being about 1.00625 (corrected).

The types bear the number 9743/10 *Zool. Surv. Ind.*

***Rhynchoplax introversus*, sp. nov.**

The carapace is ovate and is widest behind its middle point; its greatest breadth slightly exceeds its length, rostrum included. The upper surface is much sunken; in addition to the usual grooves, which are sharply demarcated, the branchial regions are traversed by a fine oblique line. There is an obscure angulation on the antero-lateral margin midway between the eye and the chelipedes, but there are no teeth in this position and no tooth or process above the base of the first walking legs. At first sight the antero-lateral and postero-lateral borders on each side appear to be discontinuous (text-fig. 11a). This, however, is due to the fact that the lateral walls of the branchial chamber project on either side and are reflected upwards, so as to form a crest which is actually higher than the true postero-lateral border. This border is continued as a low ridge within and parallel to the branchial crest. The posterior margin is short, with a slight emargination on either side opposite the last leg. The rostrum is composed of three blunt processes, the median horizontal, parallel-sided and about twice as long as broad, the two others shorter, and projecting obliquely upwards.

Almost the whole of the eye is visible from above, together with a small post-ocular tooth. The antennules fold beneath the front and are separated at the base by a blunt longitudinal ridge. The epistome is long. The external maxillipedes are of the usual form; they gape widely in the middle line and the merus is a little longer than the ischium. The exognath is almost entirely exposed (text-fig. 11b).

The chelipedes are stout and clothed with fine hairs. The merus is without teeth. The chela is stout in the male, about two and a half times as long as broad, with the palm slightly swollen. The fingers are fully one and a half times as long as the upper border of the palm; they meet throughout their length when the claw is closed and their inner margins bear five or six interlocking teeth.

The walking legs are slender; the second pair is about two and a third times as long as the carapace and rostrum. The merus in all four pairs bears a small tooth at the distal end of the upper border. The dactylus is moderately curved and is armed




FIG. 11.—*Rhynchoplax introversus*, sp. nov.
a. Large male in dorsal view. b. Anterior part of carapace of same specimen from below. c. Abdomen of same specimen.

with 6 to 8 small rather widely separated teeth that occupy the greater part of the length of the posterior border.

In the abdomen of the male (text-fig. 11c) the penultimate piece is of the usual form, but is very strongly narrowed distally. The terminal segment is exceptionally long, its length being nearly twice its basal breadth.

The species is described from two males, in one of which—much the larger of the two—the carapace is 5.4 mm. in length.

In most respects *R. introversus* is intermediate between *R. exiguus* and *R. inachoides*, but differs from both in the form of the postero-lateral border of the carapace. It resembles *R. inachoides* in the character of the dactyli of the walking legs, but the legs themselves are shorter, the rostral lobes shorter and blunter and the carapace broader.

The specimens were found by Dr. Annandale in the Tai Hu Lake in the Kiangsu Province of China. The larger individual was found off the mouth of the Tong Dong Ding Creek and the smaller at the mouth of the Moo Too Creek. Both were dredged in water about 2 metres in depth. Dr. Annandale noted that the specimens were pale buff in colour with brown markings on the carapace somewhat like a *fleur-de-lys*.

The species is remarkable in that it was obtained in pure fresh water far beyond the reach of tidal influence. A considerable number of Hymenosomatidae have been found in localities where the salinity is low and some appear to be able to exist in water that is quite fresh during a portion of the year. But the only species hitherto recorded from permanently fresh water is *Halicarcinus lacustris* (Chilton),¹ which has even been found 3,000 ft. above sea-level.

The type specimen, the larger of the two individuals, bears the number 9730/10 *Zool. Surv. Ind.*

Rhynchoplax inachoides (Alcock).

1900. *Hymenicus inachoides*, Alcock, *Fourn. Asiat. Soc. Bengal*, LXIX, p. 388, and (1902) *Illust. Zool. 'Investigator'*, pl. lxix, fig. 1.

I have little to add to Alcock's description of this species. The post-ocular denticle is clearly visible in dorsal view; the fingers of the chela of the male meet throughout their length and are armed with a regular series of teeth; the abdomen in the same sex is narrow and similar to that of *R. demeloi*.

The only known specimen is the male described by Alcock and found by Wood-Mason, along with *R. wood-masoni*, at Port Canning near Calcutta. On a recent tour in this locality I tried to obtain further specimens but was unable to find either species.

¹ For references see p. 247.

***Rhynchoplax nasalis*, sp. nov.**

The carapace is almost exactly circular and is nearly or quite as broad as long, excluding the rostrum. The surface is sunken, covered with hair, and with the grooves sharply defined. The border is entire, upturned, and is continuous from side to side across the base of the rostrum.

The rostrum differs from that of all other species in the genus in the suppression of the lateral processes; it consists merely of a single horizontal plate, more than twice as long as wide, pointed at the apex and bordered with hairs (text-fig. 12).




FIG. 12.—*Rhynchoplax nasalis*, sp. nov.

The basal segment of the antennular peduncle and the whole of the eye are visible in dorsal view. There is no post-ocular tooth and no trace of an inter-antennular septum. The epistome is of moderate length. The external maxillipedes are similar to those of the preceding species, but the merus is larger in proportion to the ischium and, when normally folded, the stalk of the exognath extends much beyond the distal end of the merus.

The chelae are swollen in both sexes and are much stouter than the walking legs; they are only a trifle larger in the male than in the female. The distal end of the lower border of the merus ends in a stout tooth, but the segment is not otherwise

armed. The carpus is smooth.

The chela is not greatly compressed and is not carinate either above or below. In the male it is little more than twice as long as deep, the dactylus being about one and a half times the length of the upper border of the palm. When the claw is closed the fingers meet throughout their length; they are armed with a regular series of six blunt




FIG. 13.—*Rhynchoplax nasalis*, sp. nov.
Chela of male (denuded).

teeth which diminish in size from behind forwards (text-fig. 13).

The second pair of walking legs is about two and a half times the length of the carapace and rostrum, the last pair about twice the length. All the segments are very slender and there is no tooth at the end of the upper border of the merus. The dactyli are curved; that of the first pair is simple, while in the remaining three pairs there is a single small recurved tooth situated some distance behind the apex (text-fig. 14).




FIG. 14.—*Rhynchoplax nasalis*, sp. nov.
Tip of dactylus of penultimate walking leg.

The abdomen of the male (text-fig. 15) is abnormally broad, the length of the two ultimate pieces being equal to the breadth of the penultimate. The lateral margin of the latter is abruptly narrowed anteriorly and bears a large and curiously formed tubercle near its distal end. The ultimate segment is broader than long, broadly rounded apically and with elevated lateral margins. In the female the abdomen is broad, but the ultimate segment is rather more triangular than in other species.




FIG. 15.—*Rhynchoplax nasalis*, sp. nov.
Abdomen of male.

The entire animal is covered with hairs, which are comparatively long on the chelipedes and legs. The specimens when

caught were covered with a dense coating of mud which was only removed with great difficulty. When denuded the crabs were ivory white in colour, the eggs of the female being reddish orange.

The length of the carapace and rostrum in an adult male is 4.4 mm., an ovigerous female is exactly the same size.

The species is described from fifteen specimens, most of which are exceedingly small. They were dredged in the Bidyadhari river near Chingrighatta on the outskirts of Calcutta in October and December 1914. They were found in very foul water which gave specific gravities of 1.0045 and 1.0060 on the two occasions on which the locality was visited.

The types bear the number 9744/10 *Zool. Surv. Ind.*

Genus *Hymenicoides*, nov.

The carapace is nearly circular in outline, sunken, with the usual grooves sharply defined and the lateral margins upturned. The rostrum is altogether absent. The epistome is of moderate length and the buccal cavern is bounded anteriorly by a sharp ridge. The external maxillipedes are slender, gaping widely in the middle line and leaving visible parts of the underlying appendages.




FIG. 16.—*Hymenicoides carteri*, gen. et sp. nov.
Anterior part of carapace, seen from below.

The merus is much longer than the ischium and is more than two and a half times as long as wide; the dactylus is styliform and of abnormal length, reaching the posterior limit of the buccal cavern when normally flexed (text-fig. 16). The chelipedes in both sexes are stouter than the legs. In the abdomen of the male the terminal segment is trilobate and the sutures of all the segments are distinct.

This genus is related to *Rhynchoplax* and *Halicarcinus*, but differs from both in the absence of the rostrum and in the great length of the dactylus of the external maxillipedes. It resembles *Rhynchoplax* in the slenderness of the basal segments of the external maxillipedes and *Halicarcinus* in having all the segments of the male abdomen distinct.

Type and only known species,—*Hymenicoides carteri*, sp. nov.

Hymenicoides carteri, sp. nov.

The carapace is almost circular, emarginate at the base of the last legs and with the posterior border short; it is broader than long in the proportion of 21 to 20. The upper surface is greatly depressed, with the grooves well defined, and is closely covered with minute hairs. The margin is entire and upturned; anteriorly and antero-laterally it forms an even curve and bears in the middle of the front a small tuft of hairs. The rostrum is entirely absent. The basal segment of the antennule and the greater part of the eye are visible in dorsal view.




FIG. 17.—*Hymenicoides carteri*, sp. nov.

At their bases the antennules are separated by a sharp forwardly directed tooth: there is no post-ocular tooth. The epistome is of moderate length. The buccal cavern is somewhat narrowed anteriorly; its lateral borders are rather strongly curved and, as in the genus *Rhynchoplax*, its anterior and posterior edges are curved inwards (text-fig. 16). On the sternum behind the bases of the external maxillipeds there is a semicircular ridge, concave anteriorly, which bears a fringe of long hairs. The curious structure of the external maxillipeds has been referred to in the generic description. There are long hairs on the inner borders and outer surface of the ischium and merus and on both inner and outer borders of the dactylus. The exognath bears a long flagellum and,

except for a small portion at the base of the stalk, is entirely concealed from view.

The chelipedes are greatly swollen in both sexes; the chelae of the male are much larger than those of the female. The outer border of the merus bears a conspicuous tooth in front of its middle point. On the inner side of the carpus there is a longitudinal ridge which is furnished with a fringe of long hairs. The chela of the male (text-fig. 18) is less than one and a half times as long as high. There are sharp keels on both upper and lower borders of the palm, the latter being continued to the tip of the immobile finger. These keels like that on the carpus are fringed with long hairs. The inner surface of the palm is convex and the outer face bears a huge protuberance (text-fig. 19), only well developed in very large males, which culminates in a




FIG. 18.—*Hymenicoides carteri*, sp. nov.
Chela of large male, external view.




FIG. 19.—*Hymenicoides carteri*, sp. nov.
Chela of large male, dorsal view.

short crest not far from the finger cleft. The fingers are stout and in large individuals meet only at the tips. The dactylus bears two large blunt teeth in its basal half and the fixed finger two smaller ones placed just behind them; nearer the tip each finger bears four or five teeth. The dactylus is fully one and a half times the length of the upper border of the palm and is obscurely ridged dorsally. The chela of the female is similar to that of the male, but is more slender and shows practically no trace of the large protuberance on the palm. Except for the fringes of hair already mentioned the chelipede bears only a few fine and scattered setae.

The second walking legs are a little longer than the first and third and are about three times the length of the carapace; the last pair is only two-thirds the length of the second. The anterior border of the merus in all four pairs terminates in a blunt tooth. The dactyli are slender and




FIG. 20.—*Hymenicoides carteri*,
sp. nov.
Dactylus of penultimate walking leg.

curved. Close to the apex each bears a large recurved tooth, behind which a number of smaller teeth are usually found. On the first pair of legs there are generally not more than one or two such teeth; on the other legs they are more numerous (text-fig. 20) and often extend from the base to the large subterminal tooth; the maximum number observed is eleven. There are fine hairs on all the segments and a fringe on the posterior margins of the propodus and dactylus. In large males the hairs on the propodus and dactylus of the first legs are very long and numerous, forming dense tufts that retain a great quantity of mud.

The sternum and abdomen are thickly beset with hairs. In the abdomen of the male (text-fig. 21) all the sutures are distinct. The lateral margins are markedly sinuous, the widest point being at the junction of the fourth and fifth segments; the ultimate segment is trilobed terminally and is much broader than the distal width of the sixth.




FIG. 21.—*Hymenicooides carteri*,
sp. nov.
Abdomen of male.

The carapace of the largest specimen, a male, is 5.7 mm. in length.

The species is described from twenty-two specimens found on the banks of the R. Hughli at Sibpur, near Calcutta, in January 1917, by Dr. Annandale and myself. They were obtained in timber bored by *Teredo* (*Xylotria dunlopi*) lying between tide-marks. The water at the time they were found was almost or quite fresh at all states of the tide, but is doubtless brackish later in the year. There are also in the collection two specimens, both small, collected by Mr. T. Southwell near Khulna in the Gangetic delta in August 1915.

With this species I have associated the name of Dr. H. G. Carter, Officiating Director of the Botanical Survey of India, to whom I am indebted for facilities for collecting at Sibpur. The types, which are from this locality, bear the number 9746/10 *Zool. Surv. Ind.*

Genus *Elamena*, Milne-Edwards.

1837. *Elamena*, Milne-Edwards, *Hist. nat. Crust.*, II, p. 33. [Not *Elamene*, Milne-Edwards, *Ann. Sci. nat., Zool.* (3), XX, p. 223 (1853); nor *Elamene*, A. Milne-Edwards, *Nouv. Arch. Mus. Paris*, IX, p. 321 (1873).]
1900. *Elamena*, Alcock, *Journ. Asiat. Soc. Bengal*, LXIX, p. 385 (not the synonymy).

The carapace is oval, triangular or polygonal, greatly depressed, and sometimes lamellar. The upper surface is flat or concave, without the usual sharp-cut grooves, and the lateral margins

may or may not be upturned. The rostrum is broadly truncate or triangular, never tridentate or trilobate. The epistome is long, sometimes very long and is separated by a ridge from the floor of the buccal cavern. The external maxillipedes are broad and completely close the buccal cavern; the ischium is longer, sometimes much longer than the merus, and the dactylus is, as usual, short. The chelipedes of the male may or may not be stouter than the walking legs. In the abdomen of the male the 3rd, 4th and 5th segments are fused and the sutures between them obliterated.

Judging from the Indian species, this genus, here described *sensu lato*, differs from all other Hymenosomatidae in the absence of the customary grooves on the upper surface of the carapace. In the character of the male abdomen it resembles *Rhynchoplax*, but differs from that genus in the form of the rostrum and external maxillipedes. Milne-Edwards' *Trigonoplax* is, at most, a subgenus of *Elamena* (*v. infra*, p. 274).

Six species of *Elamena*, *s.l.* are now known from the Indian coasts, all being represented in the Indian Museum with the exception of *E. gracilis*, Borradaile. I have not been able to satisfy myself regarding the position of this species and have, in consequence, omitted it from the following key. It is perhaps intermediate between *Elamena*, *s.s.* and *Trigonoplax*.

I. Margin of carapace upturned; rostrum with a vertical keel on its lower surface and in frontal view T-shaped; chelipedes of male greatly swollen, much stouter than legs [dactylus of walking legs apically triunguiculate] = *Elamena s.s.*

A. Carapace as broad or broader than long¹; rostrum broad and squarely truncate; a small post-ocular tooth present, but not visible from above

E. truncata
(Stimpson).

B. Carapace longer than broad¹; rostrum prominent, triangular; no post-ocular tooth

E. sindensis,
Alcock.

II. Margin of carapace not upturned; rostrum at most with a small tooth at base of lower surface, not T-shaped in frontal view; chelipedes of male slender, not stouter than walking legs. = subgen. *Trigonoplax*.

A. Rostrum parallel-sided at base; a strong post-ocular tooth visible in dorsal view; dactyli of walking legs armed in their distal third with a series of small teeth [carapace about as long as broad]

E. (T.) cimex,
Kemp.

B. Rostrum strictly triangular, its sides convergent from base to apex; no post-ocular tooth visible in dorsal view; dactyli of walking legs triunguiculate at apex.

1. Carapace longer than broad,¹ its antero-lateral margins curved and not longer than postero-lateral; rostrum flat above; a post-ocular tooth visible only from

¹ Rostrum included.

below; 2nd walking legs less than 2½ times length of carapace¹ ... *E. (T.) xavieri*, sp. nov.

2. Carapace broader than long,¹ its antero-lateral margins straight and very much longer than postero-lateral; rostrum hollowed above; no post-ocular tooth; 2nd walking legs more than 3 times length of carapace¹ ... *E. (T.) unguiformis*, de Haan.

Elamena truncata (Stimpson).

1858. *Trigonoplax truncata*, Stimpson, *Proc. Acad. Nat. Sci. Philadelphia*, X, p. 109 [55].
 1873. *Elamene truncata*, A. Milne-Edwards, *Nouv. Arch. Mus. Paris*, IX, p. 323.
 1893. *Elamene truncata*, Henderson, *Trans. Linn. Soc., Zool. (2)*, V, p. 395.
 1900. *Elamena truncata*, Alcock, *Fourn. Asiat. Soc. Bengal*, LXIX, p. 386.
 1906. *Elamena truncata*, Baker, *Trans. Roy. Soc. S. Australia*, XXX, p. 112, pl. II, figs. 2, 2a-d.
 1907. *Trigonoplax truncata*, Stimpson, *Smiths. Misc. Coll.*, XLIX, p. 146.

There does not appear to be any reason to doubt that the descriptions given by Stimpson and A. Milne-Edwards refer to the




FIG. 22.—*Elamena truncata* (Stimpson) ♀.

same species, though the specimens examined are from widely distant localities. A. Milne-Edwards, however, seems to have

¹ Rostrum included.

been unaware of the existence of Stimpson's account, for he makes no reference to it and his description is headed "*Elamene truncata* (nov. sp.)." That both authors have used the same specific name is presumably due to a remarkable coincidence.

Alcock was able to examine only a single example of this species, but two others have since been obtained; unfortunately all three specimens are females. Both Stimpson and A. Milne-Edwards note that the chelae of the male are inflated and there is consequently little doubt that the species belongs to *Elamena*, *sensu stricto*. As in *E. sindensis*, the margins of the carapace are up-turned and the front, or rostrum, bears on its underside a deep vertical keel, giving it a T-shaped appearance in facial view.

The abdomen of the male, according to Baker's description and figure, consists of five pieces, whereas only four are to be found in all the males of other species of *Elamena* and *Trigonoplax* that I have seen. From the figure it looks as if only the 3rd and 4th segments were fused in *E. truncata*, in place of the 3rd, 4th and 5th. A fresh examination of males is desirable.

In the Indian specimens the carapace is proportionately broader than in those described by Baker, the breadth being decidedly greater than the length. The front, or rostrum is squarely truncate, not rounded as described by Henderson. Behind the base of the swollen eyestalk there is a small post-ocular tooth (not shown in Baker's figure) which is altogether invisible in dorsal view. The chela of the female is little stouter than the walking legs; the fingers gape slightly when closed and are armed on their inner margins with minute teeth and short hairs. The dactylus in all four pairs of walking legs is triunguiculate at the apex. The anterior border of the ultimate segment of the abdomen of the female is strongly sinuous (text-fig. 23).

Alcock examined a single individual of this species, obtained at the Nicobars. The two additional specimens were found at Port Blair in the Andamans under a block of coral exposed at low water; the carapace of the larger is 4.8 mm. in length. When alive the carapace was brown in colour with four cream-coloured marks¹ as described by Stimpson.

There is also in the Indian Museum a female specimen of *E. truncata*, unfortunately with all the legs missing, received many years ago from the Godeffroy Museum under the name *Elamena quoyi*. It bears the label "Samoa and Viti Is."

The species appears to be one of wide Indo-pacific distribution. In addition to the above records it is known from the Ceylon coast (Henderson), the Loo Choo Is. (Stimpson), New Caledonia (A. Milne-



FIG. 23.—*Elamena truncata*
(Stimpson).
Terminal segment of abdomen of
female.

¹ Shown by dotted lines in text-fig. 22.

Edwards) and S. Australia (Baker). Lenz's record from Zanzibar¹ is erroneous, the specimens described belonging in all probability to Desmarest's *E. mathaei*.

Elamena sindensis, Alcock.

1900. *Elamena sindensis*, Alcock, *Fourn. Asiat. Soc. Bengal*, LXIX, p. 386 and (1902) *Illust. Zool. 'Investigator'*, pl. lxiv, fig. 3.

This species is still represented in the Indian Museum only by the specimens described by Alcock from Karachi. In addition to particulars noted by Alcock, it may be mentioned that the dactylus of the male chela bears a low blunt tooth near the base, the margin of both fingers being otherwise finely serrate. The dactylus of the walking legs is apically triunguiculate. The abdomen of the male is rather broadly triangular, its sides being lightly sinuous, with the 3rd, 4th and 5th somites fused. The terminal segment of the abdomen of the female resembles that of *E. truncata*.

Subgenus *Trigonoplax*, Milne-Edwards.

1853. *Trigonoplax*, Milne-Edwards, *Ann. Sci. nat., Zool.* (3), XX, p. 224.
1900. *Trigonoplax* (subgenus of *Elamena*), Alcock, *Fourn. Asiat. Soc. Bengal*, LXIX, p. 386.

I agree with Alcock that *Trigonoplax* can only be regarded as a subgenus of *Elamena*. *E. (Trigonoplax) xavieri*, which is described below, still further emphasizes the close relation between the two groups, the inter-antennular septum being a prominent plate, exactly as in *Elamena*, s.s.

In the subgenus the carapace is flatter than in *Elamena*, with its margins scarcely at all upturned, and the chelipedes are similar in the two sexes and not appreciably stouter than the walking legs. In the two species of *Elamena* that I have seen, the rostrum is T-shaped when viewed from in front, owing to the presence of a large vertical plate on its lower side; this structure, which is quite distinct from the septum between the bases of the antennules, is either absent in *Trigonoplax* or is represented by a tooth situated far behind the anterior margin.

These distinctions are slight and *Trigonoplax* in course of time will probably find a place in the synonymy of *Elamena*. Borra-daile's *E. gracilis* appears from the description and figure to be intermediate between the two groups here recognised.

Elamena (Trigonoplax) cimex, Kemp.

1915. *Elamena (Trigonoplax) cimex*, Kemp, *Mem. Ind. Mus.*, V, p. 216, text-figs. 4, 5, pl. xii, fig. 3.

The species differs from all related forms in the areolation of the carapace; the gastric, cardiac and hepatic regions are each

¹ Lenz, *Abhandl. Senck. naturf. Ges. Frankfurt*, XXVII, p. 367, pl. xlviii, figs. 15 a, b (1902).

slightly tumid and are separated by broad and shallow furrows. In this respect there is perhaps some approach to the condition found in *Halicarcinus* and *Rhynchoplax*, but there is no trace of the finely cut grooves that are conspicuous in those genera. The tooth on the lower surface of the rostrum, which is well marked in *E. (T.) xavieri* and slightly indicated in *E. (T.) unguiformis*, is in this species altogether absent. The dactyli of the walking legs bear a series of small teeth and are not apically triunguiculate as in all other Indian species of *Elamena*.

Elamena (Trigonoplax) cimex has hitherto been found only in the Chilka Lake, on the Orissa coast of the Bay of Bengal. The specimens were dredged in fresh water, but in a situation subject to great seasonal variation in salinity.




FIG. 24.—*Elamena (Trigonoplax) xavieri*, sp. nov.

***Elamena (Trigonoplax) xavieri*, sp. nov.**

The carapace closely resembles that of *E. (T.) cimex* in outline, but the antero-lateral borders are more strongly arched; its length is to its breadth as 13 to 12. There are shallow emarginations opposite the bases of the last two legs. The surface is quite flat, the regions not being defined in any way, and is altogether devoid of hairs; the margins are not upturned. The rostrum is a large triangular plate and is flat above; its margins are slightly convex and converge regularly from the base to the apex; they are not parallel at the proximal end as in *E. (T.) cimex*. On the under side,

near the base, the rostrum bears a sharp forwardly directed tooth (text-fig. 25).

The eyes and a small portion of the eyestalks extend beyond the carapace. A small post-ocular tooth may be seen when the carapace is viewed from beneath, but in dorsal view is altogether invisible. The antennules




FIG. 25.—*Elamena (T.) xavieri*, sp. nov.
Anterior part of carapace seen from below.

are separated at the base by a well-defined septum, much more distinct than in *E. (T.) unguiformis*. The epistome is long. The anterior border of the buccal cavern is convex on either side of the middle line. In the outer maxillipedes the ischium is much longer than the merus and is separated from it by a very oblique suture. The exognath bears a long flagellum and its basal part, though

largely overlapped by the endopod, is visible throughout its length.

The chelipedes are alike in the two sexes and are not appreciably stouter than the walking legs; they are about as long as the carapace and rostrum. The merus is without teeth and the merus, carpus and palm are slightly roughened and bear very fine and exceedingly short hairs. The chela is about four and a half times as long as high and the fingers are equal in length with the palm. Towards their apices the fingers are slightly inturned and on the inner face of the chela are somewhat hollowed longitudinally. When the claw is closed the fingers meet throughout their length;




FIG. 26.—*Elamena (Trigonoplax) xavieri*, sp. nov.
Chela of male.

each being provided with a series of small recurved teeth extending from the base to the apex (text-fig. 26).

The second pair of walking legs is slightly longer than the first or third, about two and a third times the length of the carapace; the fourth is much the shortest, about one and three quarter times the length of the carapace. All the segments except the dactylus

are roughened like the chelipedes and are thinly clothed with very fine hairs. In all four pairs the merus and carpus end in a strong tooth. The dactyli are slender and curved; the inner margin is densely fringed with hair and bears near the apex two stout recurved teeth, as in *E. (T.) unguiformis* (text-fig. 27).

The ultimate segment of the abdomen of the male is triangular, a little broader than long and with a pair of rather conspicuous pits near its base; proximally it is a little wider than the contracted distal end of the preceding portion. The distal margin of the abdomen of the female is slightly sinuous, as in *E. (T.) unguiformis*; in *E. (T.) cimex* it is more convex.

The carapace of a large female is 9.2 mm. in length; males are smaller, not exceeding 7.5 mm.

In living specimens the carapace is dark brown or slate-coloured, with pale antero-lateral margins and, as in *E. truncata*, a pair of elongated pale blotches project inwards and forwards from the bases of the last two pairs of legs.

In general appearance this species bears much resemblance to *E. (T.) cimex*; but it is in reality more closely allied to *E. (T.) unguiformis*. This is clearly shown by the presence of the inter-antennular septum and the tooth on the lower surface of the rostrum (both of which are in fact better defined than in *E. (T.) unguiformis*), and it is also evident in the structure of the dactylus of the walking legs.

The species is described from three males and three females obtained in the Mandavi river, opposite the town of Nova Goa in Portuguese India. They were dredged at a depth of about 10 feet on a muddy bottom in places where the current ran swiftly. The specific gravity of the water in which they were taken was very low, about 1.0010 (corrected).

In the specific name allusion is made to St. Francis Xavier, whose remains lie interred at Goa, not far from the place where the specimens were obtained.

The types bear the number 9750/10 *Zool. Surv. Ind.*

***Elamena (Trigonoplax) unguiformis*, de Haan.**

1839. *Ocypode (Elamene) unguiformis*, de Haan, in Siebold's *Fauna Japonica, Crust.*, p. 75, pl. xxix, fig. 1; pl. H.
 1900. *Elamena (Trigonoplax) unguiformis*, Alcock, *Journ. Asiat. Soc. Bengal*, LXIX, p. 387.
 1907. *Trigonoplax unguiformis*, de Man, *Trans. Linn. Soc., Zool.* (2), IX, p. 396.
 1915. *Trigonoplax unguiformis*, Parisi, *Atti Soc. Ital. Sci. nat.*, LIV, p. 281.

Other references are given by Alcock.

This well-known species differs conspicuously from the two preceding forms in the shape of the carapace, the antero-lateral




FIG. 27.—*Elamena (T.) xavieri*, sp. nov.
Tip of dactylus of last walking leg.

borders being proportionately very much longer and quite straight (text-fig. 28). The rostrum is hollowed above and bears near the proximal end of its lower surface a low ridge in place of the tooth found in *E. (T.) xavieri*. The epistome is extremely large, almost as long as the external maxillipedes. The fingers of the chelae are




FIG. 28.—*Elamena (Trigonoplax) unguiformis*. de Haan.
Outline of carapace.

furnished with minute teeth and the dactylus of the walking legs is trianguiculate. The abdomen of the male is broad at the base and narrow at the apex, the lateral margins being concave; the 3rd, 4th and 5th segments are fused.




FIG. 29.—*Elamena (T.) unguiformis*, de Haan.
Dactylus of first walking leg.

McCulloch¹ appears to be right in regarding his South Australian specimens as a distinct variety of this species. In Indian specimens the rostrum is not nearly so long, nor the dactyli of the walking legs so broad as shown in his illustration. I give here, for comparison, outline figures of the carapace and dactylus of the first leg.

The species is not uncommon at Port Blair in the Andamans, living among weeds in pure sea water at depths of 2 to 8 fathoms. In life, specimens are of a dull semitransparent brownish or greenish tint, without any conspicuous markings.

The species is known from the Gulf of Martaban (Henderson) and from numerous localities in Japan (de Haan, Ortmann, de Man, Parisi).

¹ *Trigonoplax unguiformis* var. *longirostris*, McCulloch, *Rec. Australian Mus.*, VII, p. 59, pl. xii, fig. 3 (1908).

***Elamena gracilis*, Borradaile.**

1906. *Elamena gracilis*, Borradaile, in Gardiner's *Faun. Geog. Maldive and Laccadive Archipel.*, II, p. 684, text-fig. 122 a, b.
1911. *Elamena gracilis*, Rathbun, *Trans. Linn. Soc., Zool.* (2), XIV, p. 242.

I have seen no specimens of this species and do not know whether it should be referred to *Elamena*, s.s., or to *Trigonoplax*. Judging from the figure the lateral margins of the carapace are up-turned; but the chelae are described as slender and apparently do not show any sexual differences. There is no mention of a vertical keel on the lower face of the rostrum.

In the form of the carapace *E. gracilis* differs conspicuously from any Indian species of the genus that I have seen. It was described by Borradaile from Minikoi and Male Atoll and has since been recorded by Miss Rathbun from Coetivy.